

The resident surgeon phacoemulsification learning curve at Clinical Department of Ophthalmology, Sestre milosrdnice University Hospital Center

Lacmanović Lončar, Valentina; Tadić, Rašeljka; Dujmović, Lana; Knežević, Lana; Koluder, Ana; Vidović, Marta; Krolo, Iva; Kaurić, Željko; Gašpar Mitrečić, Vanda; Petric Vicković, Ivanka; ...

Source / Izvornik: **Acta clinica Croatica**, 2016, 55, 549 - 554

Journal article, Published version

Rad u časopisu, Objavljena verzija rada (izdavačev PDF)

<https://doi.org/10.20471/acc.2016.55.04.03>

Permanent link / Trajna poveznica: <https://um.nsk.hr/um:nbn:hr:220:533622>

Rights / Prava: [In copyright](#) / [Zaštićeno autorskim pravom](#).

Download date / Datum preuzimanja: **2025-01-23**



Repository / Repozitorij:

[Repository of the Sestre milosrdnice University Hospital Center - KBCSM Repository](#)

THE RESIDENT SURGEON PHACOEMULSIFICATION LEARNING CURVE AT CLINICAL DEPARTMENT OF OPHTHALMOLOGY, SESTRE MILOSRDNICE UNIVERSITY HOSPITAL CENTER

Valentina Lacmanović Lončar¹, Rašeljka Tadić¹, Lana Dujmović¹, Lana Knežević²,
Ana Koluder², Marta Vidović³, Iva Krolo¹, Željko Kaurić⁴, Vanda Gašpar Mitrečić⁵,
Ivanka Petric Vicković¹, Renata Iveković¹ and Zoran Vatauvuk¹

¹Clinical Department of Ophthalmology and Optometry, Sestre milosrdnice University Hospital Center, Zagreb;

²Dr Tomislav Bardek General Hospital, Koprivnica; ³Nova Gradiška General Hospital, Nova Gradiška;

⁴Dr Ivo Pedišić General Hospital, Sisak; ⁵Karlovac General Hospital, Karlovac, Croatia

SUMMARY – The aim of the study was to analyze the resident learning process of phacoemulsification cataract surgery and to assess the resident phacoemulsification learning curve. This prospective study comprised 86 consecutive cases of phacoemulsification performed using standard technique by eight residents over a one-year period at Clinical Department of Ophthalmology, Sestre milosrdnice University Hospital Center in Zagreb. Operative data on each case included resident and attending surgeon, date of operation, step-by-step success questionnaire, and intraoperative complications. Postoperative data included visual acuity at one-month postoperative visit and postoperative complications. Eighty-six operations were performed under the supervision of attending surgeon. The mean best-corrected visual acuity improved from 0.47 ± 0.25 to 0.8 ± 0.25 postoperatively. Intraoperative complications occurred in 13 (15.12%) cases and continued to decrease throughout residency training. The mean number of individually performed steps *per* operation increased throughout residency (from 7.6 to 9.0). There was one early postoperative complication, endophthalmitis, which was successfully treated, and no other postoperative complications occurred. Resident surgical competency can be improved by maximizing the number of cataract procedures since the complication rates and the number of individually performed steps improved continuously with increasing surgical experience.

Key words: *Cataract surgery; Resident learning curve; Phacoemulsification; Posterior capsular tear; Intraoperative complications*

Introduction

Phacoemulsification is the gold standard for cataract extraction all over the world, as well as the most common surgical procedure. Still, it is one of the most challenging procedures to master during residency. As it has become the procedure of choice for most cata-

ract surgeries, learning phacoemulsification technique is an important part of resident education. The Accreditation Council for Graduate Medical Education requires 86 primary surgeon cases to be completed by the end of residency in the United States¹, while the Royal College of Ophthalmologists requires 50 phacoemulsification cases completed by residents over the first 2 years and 350 after 7 years of training². Unfortunately, even though ophthalmology residents in foreign countries spend a large proportion of their surgical training learning cataract surgery³, in our country it was not common practice⁴. As many previous studies

Correspondence to: *Rašeljka Tadić, MD*, Clinical Department of Ophthalmology and Optometry, Sestre milosrdnice University Hospital Center, Vinogradska c. 29, HR-10000 Zagreb, Croatia
E-mail: raseljkas@gmail.com

Received January 27, 2016, accepted April 25, 2016

showed that complication rates and outcomes of cataract surgery performed by residents were acceptable⁵⁻¹², it was decided to start first resident phacoemulsification-learning program at the Clinical Department of Ophthalmology, Sestre milosrdnice University Hospital Center. The purpose of this pilot study was to analyze the resident learning process of phacoemulsification cataract surgery at the Clinical Department of Ophthalmology, Sestre milosrdnice University Hospital Center, and to assess the resident phacoemulsification learning curve in order to implement cataract surgery training into the residency program in the future.

Material and Methods

This prospective study comprised 86 consecutive cases of phacoemulsification performed by eight residents over a one-year period at Clinical Department of Ophthalmology, Sestre milosrdnice University Hospital Center in Zagreb. During the study period, resident surgeons at the second, third and last year of residency performed all cases after approval by attending ophthalmologists. Four of eight residents had prior extracapsular cataract extraction experience and others had previous wet lab experience. Preoperatively collected data included patient age, sex, operated eye, best corrected visual acuity (BCVA) and type of cataract. Patients with other ocular comorbidities were excluded. The senior cataract surgeon was present throughout

the procedures. In all cases, the residents used standard technique under topical or local anesthesia. After clear cornea incision (knife 2.4 mm) and paracentesis, continuous curvilinear capsulorhexis was performed using forceps. After hydrodissection and hydrodelineation, the lens was phacoemulsified, usually with 'dived and conquer' technique using the Alcon Infinity Vision system phacomachine. When possible, three-piece foldable MA60AC Acrysoft intraocular lens (Alcon, Fort Worth, USA) was inserted in the posterior chamber, followed by aspiration of the remaining viscoelastic. After instillation of antibiotic (Ketocef, Pliva Hrvatska, Zagreb, Croatia) in the anterior chamber, corneal incision and paracentesis were hydrated. When needed, clear corneal incision was sutured using 10-0 suture. At the end of surgery, corticosteroid subconjunctival injection was administered. At the end of surgery, the trainees completed a questionnaire defining for each of 10 steps whether they succeeded to perform it alone or with assistance of the senior surgeon. Operative data obtained on each case included resident and attending surgeon, date of operation, step-by-step success questionnaire, intraoperative complications, and resident case number representing actual experience of the individual resident. Postoperative data collected included visual acuity at one-month postoperative visit and postoperative complications. At the end of the study, all residents completed a questionnaire rating the degree of difficulty of 10 individu-

SUBJECTIVE DIFFICULTY OF EACH STEP IN PHACOEMULSIFICATION SURGERY							
NAME:							
STEP		SCORE					
CORNEAL INCISION	very easy	1	2	3	4	5	very difficult
PARACENTESIS	very easy	1	2	3	4	5	very difficult
VISCOELASTIC INJECTION	very easy	1	2	3	4	5	very difficult
CAPSULORHEXIS	very easy	1	2	3	4	5	very difficult
HYDRODISSECTION	very easy	1	2	3	4	5	very difficult
PHACOEMULSIFICATION	very easy	1	2	3	4	5	very difficult
I/A	very easy	1	2	3	4	5	very difficult
IOL INSERTION	very easy	1	2	3	4	5	very difficult
VISCOELASTIC REMOVAL	very easy	1	2	3	4	5	very difficult
HYDRATION	very easy	1	2	3	4	5	very difficult
DEXA INJECTION	very easy	1	2	3	4	5	very difficult

Fig. 1. The questionnaire to assess subjective difficulty in performing each step.

al phacoemulsification stages on a 5-point scale from 1 (very easy) to 5 (very difficult) (Fig. 1). Statistical analysis was performed using Microsoft Excel.

Results

Eight residents performed 86 operations under the guidance of three attending surgeons with relatively even case distribution among residents (mean number of cases: 10.75 ± 2.56); 73 (84.88%) procedures were successfully performed without complications, whereas some complications occurred in 13 (15.12%) cases. The mean patient age at the time of surgery was 68 ± 7.56 years. None of the procedures had to be con-

verted to extracapsular cataract extraction (ECCE). The mean BCVA improved from 0.47 ± 0.25 to 0.8 ± 0.25 postoperatively.

In order to establish learning curve for each resident, we evaluated the number of successfully individually performed steps of operation for each resident case number during training period and identified those steps that most commonly needed assistance by senior surgeon. The mean number of individually performed steps *per operation* was 7.6 at beginning and 9.3 at the end of the study (Fig. 2). Figure 3 shows the percentage of success in performing each step in all operations.

Steps that most commonly needed assistance were capsulorhexis (44.19%, mean 19), phacoemulsification




Fig. 2. The resident learning curve: number of individually performed steps per resident case number.




Fig. 3. Percentage of successful steps in all operations.




Fig. 4. Improvement in performing most difficult steps by increasing resident experience.

(19.78%, mean 34) and cortex removal (37.00%, mean 29). When comparing success in performing these steps at the beginning with cases at the end of the study, improvement in performance was recorded throughout the period of residency training (Fig. 4).

Intraoperative complications occurred in 13 (15.12%) cases and included posterior capsular tear in 13 cases and one case of nucleus drop. Posterior capsular tear occurred in 8 cases during phacoemulsification and in 5 cases during cortex aspiration. Comparison of the resident case number recorded at the beginning (first five operations) and later in the period of training showed a decrease in the rate of intraoperative complications (8.14% *vs.* 6.98%) (Fig. 5). The rate of complications continued to decrease throughout residency training. The mean postoperative follow up period was 6 months. There was one early postoperative complication, endophthalmitis, which was successfully treated, and no other postoperative complications such as cystoid macular edema or keratopathy were observed.

According to resident subjective judgment, the most difficult stages were phacoemulsification and capsulorhexis with the highest scores (4.8 ± 0.4 and 3.6 ± 0.5), followed by cortex aspiration (3.33 ± 1.4) with completion rates of 20.93%, 55.81% and 32.56%, respectively.




Fig. 5. Decrease in the intraoperative complication rate.

The last two steps of subconjunctival corticosteroid injection and hydratization of incisions, and the first two steps of corneal incision and paracentesis were easiest to perform, yielding a nearly 100% completion rate, while hydrodissection, irrigation/aspiration and intraocular lens insertion had difficulty rates of 3.3 ± 1.2 , 2.0 ± 0.89 and 3.0 ± 1.41 , with completion rates of 72.09%, 58.14% and 55.81%, respectively (Fig. 6).

Discussion

In our study, most patients achieved very good visual acuity (0.8 ± 0.25) after resident operation, which is consistent to other literature reports^{13,14}. This can be explained by good case selection and excluding patients with other preexisting ocular comorbidities. Careful patient selection appears to be very important for overall success, especially in resident early cases^{9,11,15}.

According to our results, phacoemulsification, capsulorhexis and cortex aspiration are steps that most commonly require assistance of senior surgeon (although increasing success is recorded with the increasing number of operations, i.e. resident experience). Our data suggest that the learning curve for capsulorhexis is faster than that for cortex aspiration and phacoemulsification. For mastering these steps, residents require more successive cases over time, as shown by Robin *et al.*⁷. Baxter *et al.* report that high-frequency intensive surgical training leads to safe and efficient mastering of phacoemulsification steps with a low rate of intraoperative complications (0.66%) and high completion rate after first 6 months¹².




Fig. 6. Completion rate and mean difficulty score in each step.

In our study, the overall intraoperative complication rate (15.12%) was in the upper normal range reported elsewhere (1.8 % to 15%)⁵⁻¹². This could be explained by the fact that half of the trainees (4/8) were at the very beginning of the operating learning curve, having only wet lab experience. Although some authors report that prior experience in standard ECCE is required for safe phacoemulsification technique learning, we agree with Quillen *et al.*, Rutar *et al.* and Briszi *et al.*, who report the rate of intraoperative complications to be comparable for third-year residents with and without previous ECCE experience⁹⁻¹¹. Additionally, a relatively low mean number of operations *per* resident was the greatest limitation of our study since Randleman *et al.* showed the rate of posterior capsular tear to decrease beyond the first 200 cases and the rate of vitreous loss after the first 80 cases¹³. Our study confirmed the rate of intraoperative complications to decrease during learning process with the increasing number of operations, as previously reported^{9,10,13,14,16}. Based on the report by the Residency Review Committee of Accreditation Council for Graduate Medical Education¹, according to which the minimum number of cataract procedures performed by resident as primary surgeon should be 86, we intend to continue our training program in order to improve

our resident skills and competency. Concerning subjective difficulty in each step of the operation, residents found phacoemulsification and capsulorhexis to be the most difficult steps in the learning curve. These results are similar to those reported by Dooley and O'Brien¹⁴. Lower completion rates for the last steps, which were marked as easy in our study, could be explained by the fact that in cases where previous step needed assistance of senior surgeon, he usually would finish the operation. Our study was designed for residents to start learning the procedure from the beginning and senior surgeon would intervene in the stage where resident would have problems, often finishing the operation. Comparing subjective judgment with objective completion rate of individual steps, our study indicated that learning process should be started with easiest steps, so that the need for senior surgeon to take over early in the course of surgery would be reduced to minimum. Such a step-by-step training program has already been reported by Yulan *et al.* as a safe and effective way for residents to achieve independent phacoemulsification cataract surgery skills¹⁷. The low overall completion rate in our study (4.65%) supports the step-by-step training program to be recommended in the learning process. As a result of our study, we plan to improve the quality of the resident training program

at our Department by maximizing the number of cataract procedures performed by residents and implementing the step-by-step training model.

References

1. Accreditation Council for Graduate Medical Education [Internet]. Ophthalmology resident operative minimum requirements. Available from: http://www.acgme.org/acWebsite/RRC_240/240_MinimumsOPerativeTable.pdf.
2. Royal College of Ophthalmologists Training Committee [Internet]. Guide for Basic Specialist Training in Ophthalmology. Available from: <http://www.rcophth.ac.uk/edu-train-docs/GuifetoBST.pdf>.
3. Rowden A, Krishna R. Resident cataract surgical training in United States residency programs. *J Cataract Refract Surg.* 2002;28(12):2202-5.
4. Vatavuk Z, Vrabc R, Benčić G, Čima A, Mandić K, Mandić Z. Analysis of current ophthalmology residency curriculum. *Acta Clin Croat.* 2007;46(Suppl 1):45-50.
5. Prasad S. Phacoemulsification learning curve: experience of two junior trainee ophthalmologists. *J Cataract Refract Surg.* 1998;24(1):73-7.
6. Hennig A, Schroeder B, Kumar. Learning phacoemulsification. Results of different teaching methods. *Indian J Ophthalmol.* 2004;52(3):233-4.
7. Robin AL, Smith SD, Natchiar G, Ramakrishnan R, Srinivasan M, Raheem R, Hecht W. The initial complication rate of phacoemulsification in India. *Invest Ophthalmol Vis Sci.* 1997;38(11):2331-7.
8. Thomas R, Naveen S, Jacob A, Braganza A. Visual outcome and complications of residents learning phacoemulsification. *Indian J Ophthalmol.* 1997;45(4):215-9.
9. Rutar T, Porco TC, Naseri A. Risk factors for intraoperative complications in resident-performed phacoemulsification surgery. *Ophthalmology.* 2009;116(3):431-6. doi: 10.1016/j.ophtha.2008.10.028. Epub 2009 Jan 22.
10. Briszi A, Prahs P, Hillenkamp J, Helbig H, Herrmann W. Complication rate and risk factors for intraoperative complications in resident-performed phacoemulsification surgery. *Graefes Arch Clin Exp Ophthalmol.* 2012;250(9):1315-20.
11. Quillen DA, Phipps SJ. Visual outcomes and incidence of vitreous loss for residents performing phacoemulsification without prior planned extracapsular cataract extraction experience. *Am J Ophthalmol.* 2003;135(5):732-3.
12. Baxter JM, Lee R, Sharp JA, Foss AJ; Intensive Cataract Training Study Group. Intensive cataract training: a novel approach. *Eye (Lond).* 2013;27(6):742-6.
13. Randleman JB, Wolfe JD, Woodward M, Lynn MJ, Cherwek DH, Srivastava. The resident surgeon phacoemulsification learning curve. *Arch Ophthalmol.* 2007;125(9):1215-9.
14. Dooley IJ, O'Brien PD. Subjective difficulty of each stage of phacoemulsification cataract surgery performed by basic surgical trainees. *Cataract Refract Surg.* 2006;32(4):604-8.
15. Blomquist PH, Morales ME, Tong L, Ahn C. Risk factors for vitreous complications in resident-performed phacoemulsification surgery. *J Cataract Refract Surg.* 2012;38(2):208-14.
16. Tarbet KJ, Mamalis N, Theurer J, Jones BD, Olson RJ. Complications and results of phacoemulsification performed by residents. *J Cataract Refract Surg.* 1995;21(6):661-5.
17. Yulan W, Yaohua S, Jinhua T, Min W. Step-by-step phacoemulsification training program for ophthalmology residents. *Indian J Ophthalmol.* 2013;61(11):659-62.

Sažetak

KRIVULJA UČENJA OPERACIJE MRENE METODOM FAKOEMULZIFIKACIJE SPECIJALIZANATA NA KLINICI ZA OFTALMOLOGIJU KLINIČKOG BOLNIČKOG CENTRA SESTRE MILOSRDNICE

V. Lacmanović Lončar, R. Tadić, L. Dujmović, L. Knežević, A. Koluder, M. Vidović, I. Krolo, Ž. Kaurić, V. Gašpar Mitrečić, I. Petric Vicković, R. Iveković i Z. Vatavuk

Cilj ove studije bio je prikazati proces i krivulju učenja specijalizanata oftalmologije i optometrije u operaciji katarakte metodom fakoemulzifikacije. U ovom prospektivnom istraživanju izvedeno je 86 uzastopnih operacija katarakte metodom fakoemulzifikacije koju su u razdoblju od jedne godine na Klinici za oftalmologiju KBC "Sestre milosrdnice" učili specijalizanti oftalmologije uz nadzor iskusnih operatera. Prikupljeni operativni podaci uključili su specijalizanta operatera, mentora, datum operacije, upitnik uspješnosti po koracima operacije i intraoperacijske komplikacije. Poslijeoperacijski se pratila najbolje postignuta korigirana vidna oštrina operiranih nakon mjesec dana te razvoj poslijeoperacijskih komplikacija. Ukupno je učinjeno 86 operacija uz stalni nadzor iskusnijih operatera. Najbolje korigirana vidna oštrina operiranih poboljšala se sa srednje vrijednosti od $0,47 \pm 0,25$ na $0,8 \pm 0,25$ poslijeoperacijski. Intraoperacijske komplikacije su nastupile u 13 (15,12%) operacija, ali se njihova učestalost tijekom vremena u procesu učenja smanjila. Prosječan broj samostalno učinjenih koraka po pojedinoj operaciji za svakog specijalizanta se povećao tijekom procesa učenja (prosječno od 7,6 do 9,0). Poslijeoperacijski se samo u jednog bolesnika razvio endoftalmitis, dok druge poslijeoperacijske komplikacije nisu zabilježene. Kirurške vještine specijalizanata se mogu poboljšati povećanjem broja i učestalosti učinjenih operacija tijekom vremena, jer učestalost intraoperacijskih komplikacija opada, a uspješnost svladavanja pojedinih koraka operacije raste s povećanjem kirurškog iskustva.

Ključne riječi: *Operacija mreine; Krivulja učenja; Fakoemulzifikacija; Ruptura stražnje kapsule; Intraoperacijske komplikacije*